Connecting Deer, Forests, and Water Quality



Completed by the Students in ENVS 491
Senior Seminar, Fall 2008

Acknowledgements

We would like to thank the following people for all of their assistance and support throughout this project:

Ms. Diane Douglass, Insurance Administration, Environmental Health and Safety, Towson University

Mr. Dan Ellis, helped in the field

Mr. Bill Felter, Reservoir Natural Resources Section, Baltimore City

Mr. Rob Hirsh, GIS Analyst, Baltimore County DEPRM

Mr. Clark Howells, Reservoir Natural Resources Section, Baltimore City

Mr. Ryan Mazeska, Reservoir Natural Resources Section, Baltimore City

Mr. John Miller, helped in the field

Mr. Don Outen, Natural Resource Manager, Baltimore County DEPRM

Piney Run Golf Course Managers

Mr. Gene Scarpula, Watershed Manager, Baltimore City

Mr. Paul Schaph, helped in the field

Dr.

Table of Contents

Acknowledgements ii
Class Participants iii
Forward 2
Introduction 3
Loch Raven and Prettyboy Reservoirs 4
Current Land Use in Prettyboy and Loch Raven Reservoirs

Introduction

Loch Raven and Prettyboy reservoirs are

of pollutants is carried in runoff, and ultimately, more polluted water drains into our local reservoirs (Chesapeake Bay Program, 2008).

Locally there is now an over-abundance of white-tailed deer (Odocoileus virginianus) in the forests surrounding the reservoirs, and it has become a water quality concern in recent years. The population of deer is above what the forest can sustainably support, so white-tailed deer are now over-browsing seedlings and saplings of native forest tree13 61b4(s) 644.98 Tm 0 Tc[( )] TJ ET BTBT 1 0 0 1 2

either forested (38%) or covered by suburban and urban development (13%); 11 out of the 13% land use that is categorized as suburban/urban development is described as low-density residential land use (BC DEPRM, 2008a). An estimated 2.5% of the Prettyboy watershed consists of impervious surfaces (BC DEPRM, 2008a).

The Loch Raven Reservoir watershed is located on the Gunpowder Falls, downstream of the Prettyboy Reservoir (BC DEPRM, 2008b). The watershed for Loch Raven is located north of the Baltimore Beltway in central Baltimore County; it too drains waters from Pennsylvania in its northern most section (BC DEPRM, 2008b). Within the Maryland portion of the watershed, agriculture accounts for 37% of land use (BC DEPRM, 2008b). Therefore, Loch Raven has comparatively less agricultural land use than Prettyboy; whereas Loch Raven has 37% agricultural land use, Prettyboy has 50% agricultural land use. Another major difference between the two watersheds is that urban development accounts for 25% of the land use in Loch Raven, which is much greater than the 13% urban development in Prettyboy. Also, suburban/urban development in Loch Raven tends to be higher density than in Prettyboy. In the Loch Raven watershed, roughly 5% of the land consists of impervious surfaces (BC DEPRM, 2008b). Like Prettyboy though, forest cover accounts for 38% of the land in the Loch Raven watershed (BC DEPRM, 2008b).

Historical Land Use in Prettyboy and Loch Raven Reservoirs

It is estimated that Maryland had 95% forest cover prior to European colonization (BC DEPRM, 2008a). The original ancient forests that were once present have all but disappeared, but large areas have re-grown into mature or secondary succession forests (BC DEPRM, 2008a). These areas have re-grown from agricultural lands and from timber harvest areas (BC DEPRM, 2008a). The amount of forest cover in Prettyboy and Loch Raven, though considerably reduced by development, is greater in extent than many similarly urbanized watersheds (BC DEPRM, 2008a). Much of the forested area we see around Prettyboy and Loch Raven is protected, because of the range of benefits ±from ecological to recreational ±that these forests provide (BC DEPRM, 2008a).

The Prettyboy and Loch Raven

Forests

Role in Maintaining the Health of the Reservoirs

Forests help maintain the health of

Impervious Surfaces

Impervious surfaces contribute to excess nutrients in runoff (Chesapeake Bay Program, 2008). Impervious surfaces include roads, sidewalks, parking lots, rooftops, and any other surface which prevents water from seeping into the soil (Chesapeake Bay Program, 2008). Impervious surfaces cause less water to be absorbed into the ground. Also, the reduced plant cover means that less water is absorbed by plants too. As a result, water with greater nutrient loads can be carried off to a major water body (Chesapeake Bay Program, 2008).

Excessive sediment in runoff can cloud water, blocking sunlight from reaching submerged plants (MDE, 2008). Also, sediments settle to the bottom of streams, clogging the gravel beds used by fish for laying their eggs (MDE, 2008). Furthermore, the amount of sedimentation is an important factor in reservoir maintenance, because the accumulation of sediment reduces the amount of water that the reservoirs can hold (Ortt et al., 2000). It is estimated that between 1913 and 1997, sedimentation resulted in a lost storage capacity of 2.3 billion gallons in the Loch Raven Reservoir (Ortt et al., 2000). The annual rate of lost storage capacity for Loch Raven Reservoir is 26.8 million gallons per year (Ortt et al., 2000). Prettyboy Reservoir loses a slightly smaller amount at 23.1 million gallons of storage capacity per year (Ortt et al., 2000). Despite these staggering numbers, both reservoirs are significantly below the average for lost storage capacity in reservoirs (Ortt et al., 2000).

The problem of increased impervious surfaces leading to degraded water quality is comparatively obvious, what is less obvious is that the quality of forest cover is also a factor in water quality.

in terms of finding a suitable location for nesting (Bill McShea, qtd. in Ness,

mammals (Huebner et al., 2006). This vine also has the potential to interbreed with our native American bittersweet vine (C. scandens), creating a more fertile offspring (Huebner et al., 2006).

Mile-a-minute weed (Persicaria perfoliata), also known as Asiatic tearthumb, is an open and shade-tolerant annual vine (Huebner et al., 2006). It has a seed bank that lasts three years and is dispersed by humans, birds, and small mammals (Huebner et al., 2006).

their home ranges and the amount of seasonal movement between these areas depends greatly on the availability of food (DeNicola et al., 2000). At the local latitude (39°N) very little movement between summer and winter feeding areas is observed (DeNicola et al., 2000).

Bucks grow and shed their antlers seasonally in December or January, after mating season (DeNicola et al., 2000). Surprisingly, antlers are not often found lying around in forests; instead, they are quickly eaten by other forest creatures, which use them as a source of calcium (Michigan DNR, 2008). Antlers begin to re-grow in April in preparation for the next mating season (DeNicola et al., 2000). During mating season, also called the rut, bucks will fight with one another by charging and pushing in order to establish dominance and gain a mate (Dewey et al., 2003).

White-tailed deer breed once a year generally beginning in October and ending in December (DeNicola et al., 2000). If a doe is not impregnated during her first seasonal heat, which lasts about 24 hours, than she will go into a second heat about a month later (Dewey et al., 2003). The gestation period is six months with most fawns born in May through July (Dewey et al., 2003; DeNicola et al., 2000). The first time a doe gives birth she will only have one fawn, but in the following years she may have up to four fawns per birth (Dewey et al., 2003). At birth, fawns weigh about 4.5-5.5 lbs depending on their gender (Dewey et al., 2003). In general, young females stay with their mother longer than bucks do, though does may reach sexual maturity earlier than bucks (Dewey et al., 2003). The average age of sexual maturity for both males and females is 2 years (Dewey et al., 2003).

White-tailed deer are strict herbivores (Dewey et al., 2003). They have lower incisors, which they push against their upper pad for biting, and large molars for chewing (Fox & Myers, 2001). A twig that has been browsed by a deer is distinctive; since they do not have upper incisors, they leave an unclean, ragged tear, whereas a different animal like a rabbit would leave a clean cut (Michigan DNR, 2007). They are primarily grazers in the summer, feeding

Annuals, Perennials, and Bulbs
Asarum spp. Ginger
Aster spp. Aster
Oxalis spp. Sorrel

Kalmia latifolia Mountain Laurel
Vaccinium corymbosum Highbush Blueberry
Viburnum sp. Viburnums

Annuals, Perennials, and Bulbs
Polygonatum biflorum 6 R O R P S e q ¶ V
Ranunculus asiaticus Buttercup
Trillium spp. Trillium
Vinca minor Periwinkle

Deer and People

For the Native Americans, deer

be helpful in terms of controlling the rising deer population, the Department of Natural Resources reports the public is concerned with

carriers of Borrelia burgdorferi, the bacterium which causes Lyme disease (Allan et al., 2006). As mentioned above, fragmented forests host few to no large predators so not only deer, but also white-footed mice, flourish. More mice, coupled with more deer, result in more Lyme disease overall (Allan et al., 2006).

Lyme disease is caused by the bacterium Borrelia burgdorferi, transmitted to humans by the bite of infected blacklegged ticks (CDC, 2008). Symptoms of Lyme disease include: fever, headache, fatigue, and erythema migrans (CDC, 2008). Erythema migrans is a skin rash with a reddish edge with normal-colored skin in the middle, frequently called a "bull's eye" rash. If observed after a tick bite, it is a strong indicator of someone with Lyme disease (CDC, 2008). The symptoms usually develop anywhere from three to thirty days after the tick bite (CDC, 2008). If left untreated, infection can spread to joints, the heart, and the nervous system (CDC, 2008). The most common later problem associated with Lyme disease is arthritis, which may occur months after the tick bite (CDC, 2008). Lyme disease is diagnosed based on symptoms, physical findings (i.e. rash), and the possibility of exposure to infected ticks (CDC, 2008). Most cases of Lyme disease can be treated successfully with a few weeks of antibiotics (CDC, 2008).

There are two less common diseases associated with or carried by deer ticks: human

Warehouse, 2008). Repair fees for bumper damage to a Civic can include the following: bumper cover $180-$220, absorber $57-$72, reinforcement bar $110-$135, brackets $27-absor

How Large is Our Deer Population?

In an attempt to become more familiar with the forests and the deer management issues, we estimated the local deer population density and



Map 1: Location of sampling sites at Prettyboy and Loch Raven Reservoirs

Prior to data collection, we reviewed descriptions of deer pellets including the description of size, textures, colors, and distribution. Our sampling was started on Sept 27 and was completed by Oct 12. Different groups of students were involved in sampling over this period. Four groups of six students sampled two neighboring transects at both the Loch Raven and Prettyboy site.

Results and Discussions

The ³ W U D G Ldeer pellet count method is performed at the end of the winter. Under those conditions, pellets that are above the leaf layer are counted and the pellets are assumed to

Line 10: Omitted [major rain event]

444/2 = 888 deer/ mi

om Loch Raven site II.

Group	Date of Sampling	# Pellets Found
Tim et al.	10/4/08	9
	10/4/08	4

Within Loch Raven or Prettyboy Reservoirs we sampled two areas that were in close proximity (see map); one might expect that these neighboring areas would have similar deer density, but based on our data

the ³ O H-ff D G Should be explored. We would suggest using a single sampling team to avoid individual variation.

Deer and regeneration surveys are currently performed at the same time. We suggest that combined surveys are unnecessary and may be subject to considerable error. Regeneration surveys in winter months may lead to difficulty distinguishing between tree species as much vegetation is without leaves. Therefore, regeneration surveys throughout the wamer parts of the year may serve as a better indication of the of regeneration of the forest.

Should Baltimore City wish to demonstrate the impact deer are having on forest regeneration, enclosures that exclude deer but allow the entry of smaller animals could be used (BRNR, n.d.). Results are usually visible in about three years (BRNR, n.d.). This method would also give us information about

trails were also recorded (McCaffery, 1976). This method would not be ideal for Loch Raven because the lack of vegetation would make deer trails difficult to identify. In

Hunting has historically been allowed in Prettyboy and Liberty reservoirs, and as of 2008 limited hunting

must be used if the deer is delivered to a taxidermist or butcher (MD DNR, 2008a). Hunters that do not wish to keep the meat from the deer can donate it to organizations such as Farmers and Hunters Feeding the Hungry, where the deer is butchered and distributed to food banks, which provide meat at low costs to those in need (MD DNR, 2008e).

While hunting has proven to be a successful means of controlling the deer population in several states, several issues surrounding hunting must still be considered. First, the number of recreational hunters has been in decline, bringing into question if recreational hunters should be required to pay to hunt (Brown et al., 2000). In some studies, hunting has also been shown to modify feeding grounds for white-tailed deer, which could cause the deer population to move to another location where hunters are not present (Nixon et al., 2008). In that case it could potentially push deer out of the reservoirs and onto private land. The modification of feeding grounds and behavior can lead to more car accidents, especially in areas with high traffic roadways like Loch Raven reservoir (Sudharsan et al., 2006),.

In order to ensure the effectiveness of hunting as a deer management method, accurate deer population estimates must be made in hunted areas (DeNicola et al., 2000). Current management plans require deer population estimates in order to ensure that proper sex and age ratios are maintained (DeNicola et al., 2000). In addition to population estimates, physical condition, mortality,

c.

permits and licensing fees are also a key contributor to funding multiple Maryland DNR projects. Regulated hunting not only removes deer from the population with little cost to the public, but is also shown to alleviate costs associated with other population management

little undeveloped land to support traditional hunting (MD DNR, 2008b). Hiring sharpshooters comes at a relatively high cost to taxpayers.

other animals that may be exposed to

there are field studies in Maryland at a fenced facility where 28 does were captured, tagged with ear tags and radio transmitters, and injected with GonaCon (National Wildlife Research Center, 2004). Their reproductive behaviors were monitored for two years, and compared with unvaccinated adult does in a nearby fenced area with a similar habitat (National Wildlife Research Center, 2004).

At the Deer Research Center at Pennsylvania State University, researchers studied the effects of GonaCon specifically on the reproductive functions of bucks from 1994 to 2004 (Killian et al., 2005). One of the major affects of the vaccine was altered antler development (Killian et al., 2005). Vaccine treated males shed their antlers earlier than non-treated males and some treated males showed a lack of muscular form typically seen in males during the rutting period (Killian et al., 2005). Death rates in vaccinated males were also significantly higher than in the control group (Killian et al., 2005). Overall, the study concluded that both single and two injections of GonaCon were successful in changing reproductive functions of male white-tailed deer for many years (Killian et al., 2005). However, because of reduced antler development, and higher death rates from pulmonary disease, the vaccine is not being considered for use in bucks (Killian et al., 2005).

PZP:

Porcine Zona Pellucida (PZP) is another immunocontraceptive that has been studied in deer. It is a protein found in the eggs of pigs, and when injected it stimulates the production of antibodies that block sperm receptors on the egg, making fertilization impossible (NIST, 2000). By inhibiting the ability to become pregnant, deer are induced to have two to four additional estrous cycles per year (NIST, 2000). PZP does not enter the food chain, and shows no side effects (PNC, 2008). The EPA is currently tesJ ET BT 1 .e

reproduction, mortality, immigration, emigration, disease and weather, after 3-4 years of using PZP there has been a 20% reduction in deer at this site, (PNC, 2008).

When using fertility control agents, it is necessary to treat a large proportion of the does, 70% to 90%, to reduce population growth (DeNicolla et al, 2000). This is due to frequently low annual mortality rates for suburban deer (DeNicolla, et al, 2000). In spite of these successes, effective contraceptive programs for fertility control of free ranging wildlife are not yet fully developed (DeNicolla, et al, 2000). It is expected that a safe and cost-effective fertility control method will not be presented within the next five to ten years (DeNicolla, et al, 2000). No anti-fertility agents for wildlife are currently commercially available, and are classified as experimental drugs (Curtis, et. al, 2000). In order to test deer with drugs, a federal Investigational New Animal Drug permit and approval from a state wildlife agency are required (DeNicolla, et al, 2000).

Deterrents

Another option for curbing the impacts overabundant deer populations

Human hair is another noncommercial repellant; it is inexpensive but inconsistent in its ability to repel deer (University of Missouri Extension, 1997). Hair bags are typically hung on the outer branches of trees, or around the perimeter of an area to be protected (University of Missouri Extension, 1997). The large size of impacted area in the reservoirs makes the use of individual hang bags ineffective unless certain areas are designated for regeneration priority areas. In addition, the plants in need of protection are small and may not be able to support the bags (University of Missouri Extension, 1997). Bar soap is also used as a deterrent, and has proven to be more effective in deterrence than human hair. Bar soap needs to be reapplied to trees as they grow older and may not be useful for the smaller plants that contribute to

in

method must be done every year until the seed bank is eventually depleted (Swearingen, 2008). The removal of invasive species is most effective when two or more methods are used together (Swearingen, 2008).

According to Pannill (2000) the tree of heaven can be managed by spraying herbicide such as glyphosate on it or by cutting it down. Cutting the tree of heaven down to the ground during the months of June or July over a period of three years will eliminate the tree completely, as will uprooting the tree and removing all roots (Pannill, 2000).

Japanese barberry can be managed through chemical and manual methods (Swearingen, 2008). A chemical application can be most effective if done during the fruiting season of the shrub, this is usually during the summer months (Swearingen, 2008). The manual removal of the shrub can be done at any time but best done when the soil is moist so all roots can be removed (Swearingen, 2008).

Orientalw

Stiltgrass can also be managed by using herbicide (glyphosate) and the graminicide (sethoxydim) according to Grafton et al. (2008). The study concluded that the herbicide, glyphosate, was the most effective at killing the stiltgrass. The 1.5% concentrate of glyphosate with surfactant killed 100% of the stiltgrass (Grafton et al., 2008). Huebner (2007) suggests management of stiltgrass should be concentrated on the edges, due to its dispersal method. + X H E Q (2007) V study found that the stiltgrass seeds stay in a close vicinity of the parent plant and are not wind dispersed, therefore its main dispersal is by animals and humans. Huebner (2007) suggests with the increase in deer population, the spread of stiltgrass will also increase.

Fencing to Help Forest Regeneration

Deer population control is a long term management issue that will take time to accomplish. As a result, alternatives to restoring the forest need to be considered to prevent the forest from continuing to lose function. The major problem with forest regeneration within the reservoirs is the over-browsing of young seedlings, which inhibits forest regeneration.

A simple method for reforestation is fencing (MD DNR, 2006). This method would require establishing fenced enclosures at predetermined spots around the area that is to be reforested (MD DNR, 2006). Of the many management options available for forest regeneration, fencing is currently the most logical and economically feasible approach in light of the abundant deer population (MD DNR, 2006). Fencing off sections of a forest restricts deer from the area and allows for regeneration of plants and trees that would typically be subjected to browsing pressure (MD DNR, 2006).Fencing allows seedlings to grow and reach a height that would not be subject to browse by deer (MD DNR, 2006). After the enclosures have been allowed to regenerate fencing would be

mouth and receive a shock, which in turn deters the deer from coming back (DeNicola et al., 2000). In places subject to deep snow a highly effective, yet more affordable, combination of

All of these options have the same goal: to fund the various projects that this research is proposing. These options are easy to implement, have a wide range of applications, and are at little to no

The Maryland Department of Natural Resources (DNR) has held at least one forum about hunting in Loch Raven and Prettyboy and posted information on the DNR website. Resources may lend themselves to online presentation but an in-person public forum may prove more useful and provide a better face to the community. Involvement in the forum could be encouraged by: contacting community groups, posting on the website, contacting local news stations, sending out newsletters, etc. Other possible ways to educate the community on the problem of deer overabundance and why hunting is the most effective method and necessary are as follows:

- Host an open house and discuss the issue in the form of a nature hike
- Sending out newsletters
- Hosting school assemblies
- Host school field trips

These events could be hosted in conjunction with other groups such as Hooked on Nature, Tributary Strategy program, and the Parks and People Foundation.

While t

hunting limits based on recent population counts. Many sites would have to be sampled to reduce variation.

- The amount of variation we saw between local sites leads us to suggest that multiple sampling sites within a reservoir always be sampled to more effectively assess the size of the local population.
- Traditionally browse estimates and pellet counts are conducted at the same time which we think would lead to one or both being inaccurate. If sample of browse levels occurs after bud break so that trees can be identified it will be at a time when pellets are actively decomposing; if browse is assessed prior to bud break the plant identification is subject to a great deal more error. Therefore we suggest they be decoupled.

Suggestions

The issue of deer management, forest health and water quality is complex and critically important to the regional community. While the need to maintain the water quality of the reservoirs is obvious, the actions needed to ensure water quality are fraught with controversy. Human development has altered our local forest communities to the point that deer populations have exploded and deer now threaten to destroy, by preventing regeneration, their own habitat and the forests that protect the waters. Yet, deer are more than just too numerous forest herbivores; they have also become symbols of ³nature H and ³wild.´ Reducing their numbers by re-introducing predators, in the form of human hunters, comes into direct conflict with their popular

The specifics of how hunting is managed need to be

Invasive species management is another important issue related to forest management that needs attention. A volunteer removal program for invasive species should be established in Loch Raven and Prettyboy. Work would start at the edges of the forests and employ two or more methods of invasive species removal, such as an herbicide and manual removal.

In addition to volunteer help to conserve the reservoir forests, funding for such forest conservation projects could be raised through additional taxes on water utility bills; after all, it is the water that is being protected. People should not be penalized for using water for necessities, such as drinking, bathing, cooking, washing clothes, and washing dishes; instead, the proposed tax could target water use amounts that are in excess of the average domestic BT 1 0 0 1 235.01 593.23nts

[BC B
Wastev
http://w
20Rate

Bennett L. J., English, P. F., & McCain, R. (1940). A study of deer populations by use of pellet-group counts. Journal of Wildlife Management, 4, 398-403.

Beringer, J., Hansen, L., Demand, J., Sartwell, J., Wallendorf, M., & Mange, R. (2002). Efficacy of translocation to control urban deer in Missouri: Costs, efficiency, and outcome. Wildlife Society Bulletin, 30 (3).

[BMCs6553.7824 595.27 Tm [(B)5(M)-28(C)-25(s)-78(T)-2(w)-12(o)-20(a)4(d)]TJ ET BT 1 0 0 1 68.83 782.4 Tf 12 TL /TT0 1 Tf

Cornell University Cooperative Extension. (n.d.). Deer population facts. Retrieved September 21, from http://wildlifecontrol.info/deer/Pages/DeerPopulationFacts.aspx

Creacy, G. (2006). Deer management within suburban areas. Retrieved September 23, 2008, from http://www.tpwd.state.tx.us/publications/pwdpubs/media/pwd_bk_w7000_1197.pdf

Crissey, M. (2003). Study: Deer overpopulation may hamper forest health, diversity of tree species. Associated Press. (6)33. Retrieved November 28, 2008 from http://www.fs.fed.us/ne/newtown_square/news/NE_news/2003/articles/boston_globe_Study_Deer_overpopul.pdf

Cutter, S. L., & Renwick, W. H. (2004). Exploitation conservation preservation: A geographic perspective on natural resource use. Hoboken, NJ: John Wiley & Sons, Inc.

Cypher, B. L., Yahner, R. H., & Cypher, E. A. (1988). Seasonal food use by white-tailed deer at valley forge national historical park. In Environmental management (Vol. 12, pp. 237-242). University Park, PA: Springer New York.

Daily, G. C. (1997). 1 D W Xsemices. Societal dependence on natural ecosystems. Washington, DC: Island Press.

[DCNR] Department of Conservation and Natural Resources (2006). Deer population

Eberhardt, L., & Van Etten, R.C. (1956). Evaluation of the pellet group count as a deer census method. Journal of Wildlife Management 20(1): 70-74.

Ehrenfeld, J.G., Haggblom, M., & Kourtev, P. S. (2003). Experimental analysis of the effect of exotic and native plant species on the structure of soil microbial communties. Soil Biology and Biochemistry. 35(7): 895-905.

Environmental Protection Agency. (1998). Nonpoint pollution of surface waters with phosphorus and nitrogen. Retrieved December 4, 2008, from http://www.epa.gov/watertrain/pdf/issue3.pdf

Environmental Protection Agency. (2007). Loch Raven & Prettyboy: Decision Rationale. Retrieved October 12, 2008, from http://www.epa.gov/reg3wapd/tmdl/MD_TMDLs/LochRaven/LochRaven&PrettyboyDR.pdf

Feldhamer, G. A., Thompson, B. C., & Chapman, J. A. (2003). Wild mammals of North America. Retrieved September 21, 2008, from http://books.google.com/books?id=xQalfqP7BcC&printsec=frontcover&dq=pre+colonial+range+of+white-tail

Fergus, C. (2003). Wildlife of Virginia and Maryland and Washington, DC. Retrieved September 21, 2008, from http://books.google.com/books?id=W7UxSPd2XMAC&printsec=frontcover&dq=+history+deer+population+maryland

Fox, D., & Myers, P. (2001). Cervidae. Retrieved December 15, 2008, from http://animaldiversity.ummz.umich.edu/site/accounts/information/Cervidae.html.

Glass.net, LLC. (2008). Retrieved October 16, 2008, from http://www.glass.net/

Grafton, W. N., McGill, D. W., Pomp, J. A. (2008). Managing Japanese stiltgrass dominated communities in the ridge valley physiographic province in eastern West Virginia. Retrieved October 7, 2008 from http://nrs.fs.fed.us/pubs/gtr/gtr-p-24%20papers/53mcgill-p-24.pdf

Green, D., G. Askins & P. West. (1997). Developing urban Deer Management Plans: The Need for Public Education. Retrieved November 10, 2008 ,from http://digitalcommons.unl.edu/cgi/viewcontent.cgi?article=1014&context=ewdcc8

Gregory, S. K. (2002). Comparison of density estimators for white-tailed deer using aerial thermal infrared videography. Retrieved October 15, 2008, from http://www.afrc.uamont.edu/kissellr/GregoryThesis.pdf.

Griffin, J. R., Schwaab, E., & Peditto, P. A. (2008). Maryland annual deer report 2007-2008. Retrieved November 11, 2008, from http://www.dnr.state.md.us/wildlife/gpar/md_annual_deer_report07-08.pdf

Hall, R. L., Cheves, R.W., Jr., Baugher, T. A., Carcaterra, S. M., Davidson, W. R., Grafton, W. N., McConnell, T. R., Selders, A. W., Williams, C. E., & Workman, D. J. (1999). Deer Control in Home Gardens Publication No. 808. Retrieved October 15, 2008, from http://www.wvu.edu/~agexten/wildlife/deercont.PDF

Heys, B., & Keys, R. (2006). The potential habitat fragmentation caused by white-tailed deer (Odocoileus virginianus) and its impact on nest predation. Retrieved October 28, 2008, from http://www.cedarcreekinstitute.org/pdf/research/corn 0 0 1 -7(e)-9(da)4(r)-3(c)-5(r)>.9re 0 G [7 Tm [( )] TJ ET BT 1

Hill, H.R. (2002). The 2002 deer pellet group surveys. Wildlife Report No 3376, Michigan Department of Natural Resources. Retrieved from http://www.michigandnr.com/publications/pdfs/HuntingWildlifeHabitat/Reports/deer_02survey_pellet.pdf

Hotton, L. D. (2008). Deer hunting: An effective management tool. Retrieved September 21, 2008, from http://www.dnr.state.md.us/wildlife/deerhuntastool.asp

Hough-Goldstein, J., Gerlach Okay, J. A., & Swearingen, J. M. (2008). Plant Conservation $ O O L D Q F H ¶ V

[MD DNR] Maryland Department of Natural Resources. (2008d). History of white-tailed deer management in Maryland. Retrieved October 9, 2008, from http://www.dnr.state.md.us/wildlife/appendixa.html

[MD DNR] Maryland Department of Natural Resources. (2008e). Hunters and farmers feeding the hungry. Retrieved September 17, 2008, from http://www.tat84(e)-ne1(d.u44(s)tat84(w)-14)-9(l-9(i)511(l-9()-9(fetri)3/hetr

of

[NIST] National Institute of Standards and Technology. (2000). Deer immunocontraception at NIST. Retrieved October 20, 2008 from http://www.nist.gov/public_affairs/factsheet/deer.htm

Nix, S (2003). Red Maple Replaces Red Oak. Retrieved October 27, 2008, from About.com Web site: http://forestry.about.com/b/2003/08/07/red-maple-replaces-red-oak.htm

Nixon, C. M., Mankin, P. C., Etter, D. R., Hansen, L. P., Brewer, P. A., Chelsvig, J. E., Esker, T. L., & Sullivan, J. B. (2008). Migration behavior among female white-tailed deer in central and northern Illinois. American Midland Naturalist, 160(1), 178-190.

Ortt, R. A., Kerhin, R. T., Wells, D. & Cornwell, J. (2000). Bathymetric survey and sedimentation analysis of Loch Raven and Prettyb 0 Tc[( )

Shaw, J. S. (2004). Suburban development benefits wildlife. Retrieved January 1, 2009, from http://www.heartland.org/policybot/results.html?artId=14868

Smart Growth Online. (2008). About smart growth. Retrieved November 8, 2008, from http://www.smartgrowth.org/about/default.asp

Sudharsan, Krishnan, Riley, S. J., & Winterstein, S. R. (2006). Relationship of autumn hunting season to the frequency of deer-veh